

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

84 COPEIA

The iris is elliptical, the head flattened on top, the scales on top of head normal. Post oculars, 2; temporals, 1, followed by 2; superior labials, 8; inferior labials, 9; genials in 2 pairs, the posterior longer than the anterior. Anal plate divided. Scales in 21 rows; gastrosteges, 160; urosteges, 48 pairs. Length of body to anus, 160 mm.; length of tail, 29 mm.

The preserved specimen is colored as follows: body gray, spotted with a dull grayish brown; top of head speckled with brown; a streak of dull brown, 2 scales wide, reaching from the eye to the nape, where it widens to the width of the side of head; a large brown spot on nape; brown spots on dorsal side of body join and thus appear as streaks; spots of sides small and separate, reaching to the gastrosteges.

This specimen is now in the collection of Stanford

University.

GEORGIA H. BENTLEY, Stanford University, Calif.

AN ALBINO SPRING FROG IN WINTER

Mr. J. Daniels brought to me on January 23, a rather lively example of Rana clamitans, remarkable for its pale yellowish color and pink pupils. It was very pale yellowish over the hind portion of the body and hind legs, and the belly and throat had a decidedly pinkish tinge. The upper surface of the head was of deeper yellow. In size it appeared a little less than half that of average spring frogs in this vicinity. It was captured during the severe cold weather on January 18, under a log, in partly torpid condition, along a ditch near Woodbury, N. J. The body was nearly uniform in its pale tints and without any traces of markings.

HENRY W. FOWLER, Philadelphia, Penna.